



PUBLICATIONS

# *JEFFERSONIANA*

*Contributions from the  
Virginia Museum of Natural History*

---

Number 19

10 January 2009

Unusual Cambrian Thrombolites from the  
Boxley Blue Ridge Quarry, Bedford County, Virginia

Alton C. Dooley, Jr.

ISSN 1061-1878

Virginia Museum of Natural History  
Scientific Publications Series

The Virginia Museum of Natural History produces five scientific publication series, with each issue published as suitable material becomes available and each numbered consecutively within its series. Topics consist of original research conducted by museum staff or affiliated investigators based on the museum's collections or on subjects relevant to the museum's areas of interest. All are distributed to other museums and libraries through our exchange program and are available for purchase by individual consumers.

Memoirs are typically larger productions: individual monographs on a single subject such as a regional survey or comprehensive treatment of an entire group. The standardized format is an 8.5 x 11 inch page with two columns.

*Jeffersoniana* is an outlet for relatively short studies treating a single subject, allowing for expeditious publication. The standardized format is a single column on a 6 x 9 inch page.

Guidebooks are publications, often semi-popular, designed to assist readers on a particular subject in a particular region. They may be produced to accompany members of an excursion or may serve as a field guide for a specific geographic area.

Special Publications consist of unique contributions, usually book length, either single-subject or the proceedings of a symposium or multi-disciplinary project in which the papers reflect a common theme. Appearance and format are customized to accommodate specific needs; page size and layout varies accordingly.

*The Insects of Virginia* is a series of bulletins emphasizing identification, distribution, and biology of individual taxa (usually a family) of insects as represented in the Virginia fauna. Originally produced at VPI & SU in a 6 x 9 inch page size, the series was adopted by VMNH in 1993 and issued in a redesigned 8.5 x 11 inch, double column format.

Jeffersoniana, Number 19, pp. 1-14  
Virginia Museum of Natural History

## Unusual Cambrian Thrombolites from the Boxley Blue Ridge Quarry, Bedford County, Virginia

ALTON C. DOOLEY, JR.  
Virginia Museum of Natural History  
21 Starling Avenue  
Martinsville, Virginia 24112, USA  
alton.dooley@vmnh.virginia.gov

### ABSTRACT

Three unusual thrombolites were collected in June 2008 from the Late Cambrian Conococheague Formation at the Boxley Materials Blue Ridge Quarry in Bedford County, Virginia. These specimens are isolated low domes with a thrombolitic core and a pustulate, stromatolitic outer layer. The two largest domes have a distinctive thickened rim around their margins. There are apparent traces across the upper surfaces of the domes that may indicate grazing by invertebrates.

The overall structure and morphology of the Boxley specimens is reminiscent of modern thrombolites forming in Lake Thetis, a saline lake in southwestern Australia. The low domes and thickened rims in Lake Thetis specimens seem to be a result of growth in a protected setting, with shallowing water levels. Based on the similarities with the Lake Thetis specimens, the Boxley thrombolites may have formed in a protected lagoonal setting with gradually dropping water levels, followed by relatively rapid inundation and burial.

### INTRODUCTION


Stromatolites are a common occurrence globally in Proterozoic and Cambrian sediments, and occur less frequently in post-Cambrian deposits up to the present day. Examination of modern forms has demonstrated that stromatolitic and thrombolitic mounds are constructed primarily by a variety of blue-green algal communities (Black, 1933; Logan, 1961; Sharp, 1969; Reid et al., 2000), although laminated structures reminiscent of algal stromatolites can be formed by other organisms or by abiotic processes (Logan et al., 1964; Ahr, 1971; Walter, 1976; Grotzinger and Rothman, 1996). Stromatolitic structures can assume a variety of morphologies

that are influenced both by the taxa present and environmental conditions (Black, 1933; Logan et al., 1961; Dill et al., 1986), and most of these have been observed in both modern and ancient stromatolites (Logan, 1961; Hoffman, 1976).

Modern stromatolites are largely restricted to hypersaline environments, with a few exceptions (Dill, 1986). The extensive geographic range of stromatolites in the Proterozoic and Cambrian suggests that stromatolites previously lived in a greater variety of habitats than do modern examples, and that the presence of stromatolites is not necessarily an indicator of an intertidal setting (Aitke, 1967; Playford and Cockbain, 1969). The decline of stromatolites beginning in the late Cambrian has led to suggestions that the rise of grazing and burrowing metazoans in the Cambrian resulted in the restriction of stromatolites to marginal environments after that time (Awramik, 1971; Walter and Heys, 1985), although open-marine stromatolites have been found in Devonian deposits (Playford and Cockbain, 1969). The presence of burrowing metazoans may have also led to the appearance of thrombolites (unlaminated algal mounds) in the Cambrian (Walter and Heys, 1985).

## GEOLOGIC SETTING

The Late Cambrian Conococheague Formation in Virginia and Maryland is a thrombolite- and stromatolite-rich limestone that has been interpreted as a shallow subtidal to peritidal deposit (Demicco, 1983; Osleger and Read, 1991). Regressive cycles up to 10 m thick tend to have basal thrombolitic beds overlain by cross-stratified oolitic grainstones grading into bioturbated and mud-cracked rhythmites (“ribbon rock” of Demicco, 1983). The cycles are capped by laminated and stromatolitic limestones and dolostones. A section of the Conococheague is exposed in the Boxley Materials Blue Ridge Quarry in Bedford County, Virginia (Demicco, 1982). This quarry is located within the Blue Ridge fault zone, and a highly deformed allochthonous block of the Conococheague Formation, overlain by the overthrust Elbrook Formation, has been exposed during quarrying operations over the last century (Figure 1). The lithologies and cycles present in the Blue Ridge Quarry are comparable to other Conococheague localities in Virginia and Maryland.




**Figure 1.** Section of the Conococheague Formation at the Blue Ridge Quarry. The black line is approximately 2 meters, and indicates one of the thrombolitic beds. Inset: map of Bedford County, Virginia showing the location of the Blue Ridge Quarry.


## DESCRIPTION

In June 2008, Boxley employee Richard Bengé discovered a large, isolated thrombolite in blast-produced tailings during operations at the Blue Ridge Quarry (VMNH 160000; Figure 2). After discovery of this specimen, additional examination of tailings from that blast revealed two additional, much smaller bioherms (VMNH 160001; Figure 3). The largest of these specimens is a low dome, subcircular in plan view with a long axis diameter of 190 cm and a short axis diameter of 180 cm. While much larger bioherms have been described (see for example Ahr, 1971), the Boxley specimen represents one of the larger complete and intact specimens collected, and is unusual among Conococheague thrombolites in that it represents a complete bioherm.





**Figure 2.** VMNH 160000 thrombolitic bioherm. A, surface view. B, profile view. Scale = 50 cm.




**Figure 3.** VMNH 160001 thrombolitic bioherms. A, surface view. B, cross-section. Arrow indicates thickened rim. Scale = 10 cm.

The internal wackestone - to - mudstone fabric with *Renalcis* algal structures of the Conococheague thrombolites was described in detail by Demicco (1982). Internally, all the Boxley specimens have a mottled appearance; throughout most of the structure laminations are sparse, discontinuous, and located only near the tops of the domes. The outer ~5 mm of the domes are stromatolitic, with distinct knobs covering the upper surface of the dome (Figure 4). These knobs are generally circular to oval in outline, and are up to approximately 3 cm in diameter.


*Renalcis* are not obvious in these thrombolites, but filaments that could represent algal structures are visible in places in the stromatolitic layer (Figure 5).

While the internal structure of the Boxley specimens is similar to that of typical Conococheague stromatolites/thrombolites, the overall morphology is quite different. Most Conococheague bioherms are widening-upward structures, which are flat-topped and merged together at their tops, often reaching thicknesses of many meters (Demico, 1982; Osleger and Read, 1991; Figure 1). The thrombolitic units are very noticeable in most of the deposits in the Blue Ridge Quarry. In contrast, VMNH 160000 and 160001 are low-profile domes that narrow upward. The two small domes are coalesced only at their bases, while the largest specimen is isolated.



**Figure 4.** Knob on the surface of VMNH 160000, showing stromatolitic layers.

A unique feature of the Boxley thrombolites is the presence of a thickened rim around the margin of the mounds. This rim completely encircles the mound in VMNH 160000 and at least partially surrounds



**Figure 5.** Scanning electron micrograph of a section of the stromatolitic layers of VMNH 160000, showing a possible algal filament. Scale = 10  $\mu$ m.

the larger mound in VMNH 160001 (Figures 2 and 3); the smallest dome seems to lack this rim. In VMNH 160000, the thickened rim is thrombolitic with a pustulate, stromatolitic outer layer, just as in the main part of the dome.

The upper surfaces of the thrombolites exhibit a small number of apparent traces, in the form of meandering grooves (Figure 6). These grooves are approximately 3 mm wide and cut across both the stromatolitic knobs and the troughs between adjacent knobs. As each trace is restricted to the stromatolitic layers (the living part of the dome), and runs tangential to the surface of the thrombolite, they are more consistent with feeding traces than residence burrows. The large, primitive chiton *Matthevia* has been interpreted as a grazer of stromatolites and is known from the Conococheague Formation (Runnegar et al., 1975), although it has not been identified at the Blue Ridge Quarry. *Matthevia* would be expected to leave a feeding trace approximately 3 mm wide, assuming that the relative proportions of its mouth and valves were comparable to those of modern chitons.





**Figure 6.** Possible feeding trace on the surface of VMNH 160000.

## DISCUSSION

In general, the Conococheague thrombolite units have been interpreted as forming in the shallow subtidal zone (Demicco, 1982, 1983), consistent with general suggestions about thrombolite distributions suggested by Aitken (1967). These thrombolitic beds are often several meters thick, with the individual thrombolitic heads coalescing into a contiguous mass at the top of the unit. This pattern is typical throughout the Conococheague, including at the Blue Ridge Quarry. However, VMNH 160000 and 160001 differ from this pattern, in that the heads are low in profile and isolated, and have unusual thickened rims. In the section of the quarry where these specimens were collected there is no obvious thrombolitic unit, and no comparable isolated thrombolites were found in place. This suggests a somewhat different history for the Boxley specimens, compared to typical Conococheague thrombolites.

The overall morphology of the Boxley thrombolites is reminiscent of modern stromatolites/thrombolites forming in Lake Thetis, a saline coastal lake in Australia (Grey et al., 1990). The morphology of the Lake Thetis thrombolites appears to be a result of unique conditions found in that environment.


Lake Thetis is a small saline lake fed by rainfall and groundwater input, with a maximum depth of about 2.25 m (Grey et al., 1990). There is a complex and diverse array of algal structures present in and around the lake, including lithified stromatolites along the southwest shore that were described in detail by Grey et al. (1990). These stromatolites are low steep-sided domes comparable in size to VMNH 160000, with diameters up to around 1.5 meters, and extend well above the lake surface (Figure 7). The centers of these domes are eroded and often collapsed, with active growth only occurring along the wetted sides of the domes. The inside of the domes are thrombolitic, with a stromatolitic outer layer. The stromatolitic layer is composed of laminated columns about 1.5 cm in diameter and approximately 5 to 15 cm thick, which give the outer surface of the dome a pustulate appearance. This is very similar to the structure of the Boxley thrombolites, although in the Boxley specimens the outer stromatolitic layer is only a few millimeters thick. The Lake Thetis domes, and particularly the stromatolitic layers, are limited to the south shore of



**Figure 7.** Thrombolitic domes from Lake Thetis, southwestern Australia. Individual domes are approximately 1 meter in diameter. Image by Ruth Ellison, retrieved on 18 November 2008 from <http://flickr.com/photos/laruth/153584043/>. Used with permission.

the lake where wave activity is low, possibly because a gelatinous coating does not develop under high-energy conditions (Grey et al., 1990).

Some of the Lake Thetis thrombolites have a thickened rim reminiscent of the Boxley specimens (Figure 7). In modern structures this rim seems to have formed as a result of the drop in water level in Lake Thetis over the last several thousand years (Grey et al., 1990; Figure 8). The exposed stromatolitic domes erode and collapse when desiccated, even as growth continues along the wetted margins of the dome. This suggests that a drop in relative water level exposing the center of the dome is necessary for the formation of the thickened rim.



**Figure 8.** Google Earth image of Lake Thetis. Black arrows indicate past shorelines of receding lake. White arrow indicates the position of modern thrombolitic domes. Image retrieved on 18 November 2008.

The Boxley thrombolites differ from the Lake Thetis examples in that the thickened rim is combined with a large, well-developed thrombolitic

dome, suggesting a more complex history of water level variation than the steady drop observed in Lake Thetis. The unique morphology can be explained by a scenario in which stromatolite growth was initiated in a protected setting, such as a back-barrier lagoon. The shallow, low-energy water would prevent the development of large bioherms while allowing the growth of broad, low domes, with simultaneous deposition of carbonate muds. A slight drop in water level restricted growth to the margins of the dome, resulting in the development of the thickened rim. A subsequent increase in water level (perhaps due to a breach of the isolating barrier) buried the largest thrombolites under coarser-grained carbonates. Efforts to locate isolated domes in situ have so far been unsuccessful, even though their approximate position is known. That this section of the quarry also seems to lack the basal thrombolitic beds typically found in Conococheague cycles suggests that this limited area experienced somewhat different conditions from the rest of the formation.

#### ACKNOWLEDGMENTS

I would like to thank Boxley Materials Company, and particularly Ab Boxley, Bill Hamlin and Tom Roller, for their donation of these specimens to the Virginia Museum of Natural History and their continued access to the Blue Ridge Quarry. Ruth Ellison provided images of the Lake Thetis stromatolites. I would also like to thank James Beard, Brett Dooley, and Andrew Moore for helpful discussions, Mary Catherine Santoro for assistance in obtaining references, Mary Carmen for SEM images, and the Jeffersoniana editorial board and outside reviewers for helpful comments.

#### REFERENCES CITED

- Ahr, W. M., 1971. Paleoenvironment, algal structures, and fossil algae in the Upper Cambrian of central Texas. *Journal of Sedimentary Petrology*, 41:205-216.
- Aitke, J. D., 1967. Classification and environmental significance of cryptalgal limestones and dolomites with illustrations from the Cambrian and Ordovician of southwestern Alberta. *Journal of Sedimentary Petrology*, 37:1163-1178.
- Awramik, S. M., 1971. Precambrian columnar stromatolite diversity: reflection of metazoan appearance. *Science* 174:825-827.
- Black, M., 1933. The algal sediments of Andros Island, Bahamas. *Philosophical*

- Transactions of the Royal Society of London, Series B., 222:165-192, Plates 21-22.
- Demicco, R. V., 1982. Upper Cambrian Conococheague Limestone, in P. T. Lyttle (ed.), Central Appalachian Geology. Geological Society of America Northeast-Southeast Joint Section Field Trip Guidebook, American Geological Institute, pp. 217-254.
- Demicco, R. V., 1983. Wavy and lenticular-bedded carbonate ribbon rocks of the Upper Cambrian Conococheague Limestone, central Appalachians. *Journal of Sedimentary Petrology* 53:1121-1132.
- Dill, R. F., E. A. Shinn, A. T. Jones, K. Kelly, and R. P. Steinen, 1986. Giant subtidal stromatolites forming in normal salinity waters. *Nature* 324:55-58.
- Grey, K., L. S. Moore, R. V. Burne, B. K. Pierson, and J. Bauld, 1990. Lake Thetis, Western Australia: an example of saline lake sedimentation dominated by benthic microbial processes. *Australian Journal of Marine and Freshwater Research* 41:275-300.
- Grotzinger, J. P. and D. H. Rothman, 1996. An abiotic model for stromatolite morphogenesis. *Nature* 383:423-425.
- Hoffman, P., 1976. Environmental diversity of middle Precambrian stromatolites, in M. R. Walter (ed.), *Stromatolites*. Elsevier, Amsterdam, pp. 599-611.
- Logan, B. W., 1961. Cryptozoon and associated stromatolites from the Recent, Shark Bay, Western Australia. *Journal of Geology* 69:517-533.
- Logan, B. W., R. Rezak, and R. N. Ginsburg, 1964. Classification and environmental significance of algal stromatolites. *Journal of Geology* 72:68-83.
- Osleger, D. and J. F. Read, 1991. Relation of eustasy to stacking patterns of meter-scale carbonate cycles, Late Cambrian, U.S.A. *Journal of Sedimentary Petrology* 61:1225-1252.
- Playford, P. E. and A. E. Cockbain, 1969. Algal stromatolites: deepwater forms in the Devonian of western Australia. *Science* 165:1008-1010.
- Reid, R. P., P. T. Visscher, A. W. Decho, J. F. Stolz, B. M. Bebout, C. Dupraz, I. G. MacIntyre, H. W. Paerl, J. L. Pinckney, L. Prufert-Bebout, T. F. Steppe, and D. J. DesMarais, 2000. The role of microbes in accretion, lamination and early lithification of modern marine stromatolites. *Nature* 406:989-992.
- Runnegar, B., J. Pojeta, Jr., M. E. Taylor, and D. Collins, 1975. New species of the Cambrian and Ordovician chitons *Matthevia* and *Chelodes* from Wisconsin and Queensland: evidence for the early history of polyplacophoran mollusks. *Journal of Paleontology* 53:1374-1394.
- Sharp, J. H., 1969. Blue-green algae and carbonates—*Schizothrix calcicola* and algal stromatolites from Bermuda. *Limnology and Oceanography* 14:568-578.

- Walter, M. R., 1976. Geysirites of Yellowstone National Park: an example of abiogenic "stromatolites", in M. R. Walter (ed.), *Stromatolites*. Elsevier, Amsterdam, pp. 87-112.
- Walter, M. R. and G. R. Heys, 1985. Links between the rise of the Metazoa and the decline of stromatolites. *Precambrian Research* 29:149-174.





### *Parts published to date*

- 1 On the taxonomy of the milliped genera *Pseudojulus* Bollman, 1887, and *Georgiulus*, gen. nov., of southeastern United States. Richard L. Hoffman. Pp. 1-19, figs. 1-22. 1992. \$2.00
2. A striking new genus and species of bryocorine plant bug (Heteroptera: Miridae) from eastern North America. Thomas J. Henry. Pp. 1-9, figs. 1-9. 1993. \$1.00.
3. The American species of *Escaryus*, a genus of Holarctic centipeds (Geophilo-morpha: Schendylidae). Luis A. Pereira & Richard L. Hoffman. Pp. 1-72, figs. 1-154, maps 1-3. 1993. \$7.00
4. A new species of *Puto* and a preliminary analysis of the phylogenetic position of the *Puto* Group within the Coccoidea (Homoptera: Pseudococcidae). Douglass R. Miller & Gary L. Miller. Pp. 1-35, figs. 1-7. 1993. \$4.00.
5. *Cambarus* (*Cambarus*) *angularis*, a new crayfish (Decapoda: Cambaridae) from the Tennessee River Basin of northeastern Tennessee and Virginia. Horton H. Hobbs, Jr., & Raymond W. Bouchard. Pp. 1-13, figs. 1a-1n. 1994. \$2.00.
6. Three unusual new epigaeic species of *Kleptochthonius* (Pseudoscorpionida: Chthoniidae). William B. Muchmore. Pp. 1-13, figs. 1-9. 1994. \$1.50.
7. A new dinosauriform ichnogenus from the Triassic of Virginia. Nicholas C. Fraser & Paul E. Olsen. Pp. 1-17, figs. 1-3. 1996. \$2.00.
8. "Double-headed" ribs in a Miocene whale. Alton C. Dooley, Jr. Pp. 1-8, figs. 1-5. 2000. \$1.00.
9. An outline of the pre-Clovis Archeology of SV-2, Saltville, Virginia, with special attention to a bone tool dated 14,510 yr BP. Jerry N. McDonald. Pp. 1-60, figs. 1-19. 2000. \$3.00.
10. First confirmed New World record of *Apocyclops dengizicus* (Lepishkin), with a key to the species of *Apocyclops* in North America and the Caribbean region (Crustacea: Copepoda: Cyclopidae). Janet W. Reid, Robert Hamilton, & Richard M. Duffield. Pp. 1-23, figs. 1-3. 2002. \$2.50
11. A review of the eastern North American Squalodontidae (Mammalia: Cetacea). Alton C. Dooley, Jr. Pp. 1-26, figs. 1-6. 2003. \$2.50.
12. New records and new species of the genus *Diacyclops* (Crustacea: Copepoda) from subterranean habitats in southern Indiana, U.S.A. Janet W. Reid. Pp. 1-65, figs. 1-22. 2004. \$6.50.
13. *Acroneuria yuchi* (Plecoptera: Perlidae), a new stonefly from Virginia, U.S.A. Bill P. Stark & B. C. Kondratieff. Pp. 1-6, figs. 1-6. 2004. \$0.60.
14. A new species of woodland salamander of the *Plethodon cinereus* Group from the Blue Ridge Mountains of Virginia. Richard Highton. Pp. 1-22. 2005. \$2.50.
15. Additional drepanosaur elements from the Triassic infills of Cromhall Quarry, England. Nicholas C. Fraser & S. Renesto. Pp. 1-16, figs. 1-9. 2005. \$1.50.
16. A Miocene cetacean vertebra showing partially healed compression fracture, the result of convulsions or failed predation by the giant white shark, *Carcharodon megalodon*. Stephen J. Godfrey & Jeremy Altmann. Pp. 1-12. 2005. \$1.50.
17. A new *Crataegus*-feeding plant bug of the genus *Neolygus* from the eastern United States (Hemiptera: Heteroptera: Miridae). Thomas J. Henry. Pp. 1-10. \$1.50.
18. Barstovian (middle Miocene) Land Mammals from the Carmel Church Quarry, Caroline Co., Virginia. Alton C. Dooley, Jr. Pp. 1-17. \$2.00.
19. Unusual Cambrian Thrombolites from the Boxley Blue Ridge Quarry, Bedford County, Virginia. Alton C. Dooley, Jr. Pp 1-12, figs. 1-8, 2009. \$ 3.00.



*Virginia Museum of*  
**NATURAL HISTORY**

---

PUBLICATIONS

21 Starling Avenue  
Martinsville, VA 24112